



**Company Name:** FREEZONE PUBLIC HEALTH PTY LTD  
**Product Name:** Ratshot One Shot Rodenticide Pellets  
**APVMA Approval No:** 80029/100069

<b>Label Name:</b>	Ratshot One Shot Rodenticide Pellets
--------------------	--------------------------------------

<b>Signal Headings:</b>	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
-------------------------	---

<b>Constituent Statements:</b>	0.05 g/kg BRODIFACOUM Contains Denatonium Benzoate (Bittering agent)
--------------------------------	---

<b>Mode of Action:</b>	
------------------------	--

<b>Statement of Claims:</b>	READY TO USE BAIT FOR USE IN COMMERCIAL, AGRICULTURAL AND DOMESTIC BUILDINGS  KILLS RATS & MICE
-----------------------------	---

<b>Net Contents:</b>	250 g - 20 kg
----------------------	---------------

<b>Restrains:</b>	DO NOT exceed 3 m between bait stations for mice or 9 m for rats. DO NOT place bait stations in open. DO NOT apply bait to crops.
-------------------	---

<b>Directions for Use:</b>	<b>Baiting Strategy:</b> Eliminate, as far as practicable, all other food sources. Bait the infested area. Place bait stations under cover. Inspect bait stations daily. Replace eaten baits. If quickly eaten, increase number of bait stations. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers. <b>SITUATION:</b> In and around agricultural, commercial and domestic buildings.  NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.
----------------------------	--

<b>Other Limitations:</b>	THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.
<b>Withholding Periods:</b>	
<b>Trade Advice:</b>	
<b>General Instructions:</b>	<p>Pellets are not subject to wind dispersal. Most deaths occur 4 to 7 days after consumption of bait. Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Record the number and locations of bait stations for purpose of replenishment and later removal.</p> <p>Mice require several bait stations, 50 g per station (less than 1 g will kill a mouse). Rats require large quantities of bait in one or two locations, 50-100 g per station (4-6 g will kill a rat). For effective control continue until feeding ceases.</p>
<b>Resistance Warning:</b>	
<b>Precautions:</b>	<p><b>PRECAUTION: KEEP OUT OF REACH OF CHILDREN</b></p> <p>Keep away from food and water and surfaces which come in contact with food and water.</p>
<b>Protections:</b>	<p><b>PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</b></p> <p>DO NOT contaminate dams, streams, rivers, waterways and drains with the bait or its used container. The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife and care should be exercised in its use. DO NOT place baits in locations which are accessible to domestic animals, livestock or birds. The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces e.g. drains. Search for and burn or bury all rodent bodies, do not place in refuse bins or rubbish tips. Ensure uneaten baits are removed and disposed of with empty containers when rodent activity has ceased.</p>
<b>Storage and Disposal:</b>	<p>Store in a tightly closed, original container in a dry well ventilated area, as cool as possible out of direct sunlight. Shake empty container onto baiting site. Break, crush or puncture and bury empty container in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Search for and burn or bury all rodent bodies, do not place in refuse bins or in rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when rodent activity has ceased.</p>
<b>Safety Directions:</b>	<p>Poisonous if swallowed. DO NOT inhale dust. DO NOT touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water. Vitamin K1 (phytomenadione) is antidotal.</p>
<b>First Aid Instructions:</b>	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Vitamin K1 (Phytomenadione) is antidotal.</p>

<b>First Aid Warnings:</b>	
----------------------------	--