

Product Name: COCKYKILL COCKROACH GEL BAIT
APVMA Approval No: 86120/114722



Label Name:	COCKYKILL COCKROACH GEL BAIT
Signal Headings:	READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	0.5g/kg Fipronil
Statement of Claims:	A crevice, crack, or spot treatment for the control of cockroaches in commercial, industrial and residential premises as per directions for use.
Net Contents:	35g; 15g; 60g
Restraints:	Do NOT place in areas that are routinely washed
How to Use:	This section contains file attachment.
Withholding Periods:	
How to Prepare:	GENERAL INSTRUCTIONS To apply the gel, remove the cap on the nozzle and touch the tip to the surface to be treated and depress plunger slightly. Recap the dispenser after treatment is completed and return cartridge to the outer pack. Reapply according to remaining level of infestation, when bait is no longer visibly present. A visual inspection of gel placements is recommended one month after initial treatment.
Precautions:	
Protections:	PROTECTION OF LIVESTOCK, WILDLIFE, AND OTHERS

Do NOT contaminate streams, rivers or waterways with the chemical or used containers.
Do NOT apply this product in any areas easily accessible to children or animals.

Storage and
Disposal:

STORAGE AND DISPOSAL Store in the closed, original container in a cool, well ventilated locked place out of the reach of children. Do NOT store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in the garbage.

Safety Directions:

SAFETY DIRECTIONS May irritate the eyes and skin. Avoid contact with eyes and skin.
Wash hands after use

First Aid Instructions:

FIRST AID If poisoning occurs, contact a doctor or Poisons Information Centre. Telephone 131126 Australia-wide.

First Aid Warnings:

DIRECTIONS FOR USE

RESTRAINT: Do NOT place in areas that are routinely washed.

SITUATION	PEST	RATE	CRITICAL COMMENTS
Domestic, Commercial and Public Service Buildings	German and other common cockroaches	Apply 1-3 spots per m ² . Recommended spot size is 0.03-0.06g. Increase the number of spots (up to 3) and spot size depending upon severity of infestation and species present.	Gel placement should be at or near areas where cockroaches gather, such as in cracks and crevices

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.